SYMPTOM IDENTIFICATION FOR AGITATION ASSOCIATED WITH DEMENTIA DUE TO ALZHEIMER'S DISEASE

Use this resource to help with:

- **Identifying and documenting agitation symptoms** or behaviors observed in residents with Alzheimer's dementia
- **Capturing observations** for communication to designated personnel for evaluation
- Asking the interdisciplinary team to determine if further assessment should be completed

The International Psychogeriatric Association uses 4 criteria to define agitation:

- The resident must meet the criteria for a cognitive impairment or dementia syndrome
- The behavior(s) must produce excess disability beyond that due to cognitive impairment* and significantly impair ≥1 of the following:
 - Interpersonal relationships
 - Other aspects of social functioning
 - Ability to perform or participate in daily living activities

- The resident must exhibit 1 of the following behaviors:
 - ≥1 agitation behavior(s) associated with emotional distress, and the behavior(s) must be recurrent for a minimum of 2 weeks. OR
 - Behavior that represents a dramatic change in the resident's usual behavior[†]
- The agitation cannot be attributable solely to:
 - Another psychiatric disorder
 - Suboptimal care conditions
 - A medical condition
 - The physiological effects of a substance

†In special circumstances, the ability to document the behaviors over 2 weeks may not be possible, and other terms of persistence and severity may be needed to capture the syndrome beyond a single episode.



^{*}According to the clinician's opinion.

Identifying the symptoms of Agitation associated with dementia due to Alzheimer's Disease

What are agitated symptoms?

Agitated behaviors are defined by the Cohen-Mansfield Agitation Inventory (CMAI). The CMAI is a clinically widespread tool that provides 29 symptoms of agitation to look for when caring for residents with Alzheimer's disease. Symptoms can be verbal, physically non-aggressive, or aggressive, and are scored by clinicians based on caregiver input.

Resident name							
	Check any symptom(s) observed Agitation behaviors as defined by CMAI						
Aggressive	Physically Non-aggressive	(Dea)	Verbally Agitated				
Screaming	Pacing, aimless wandering		Complaining				
Biting	General restlessness		Constant unwarranted request for attention or help				
Hitting	Inappropriate dress or disrobing	П	Repetitive sentences or question				
Kicking	Trying to get to a different place		Negativism				
Hurting self or others	Handling things inappropriately						
Cursing or verbal aggression	Performing repetitive mannerisms						
Pushing							
Scratching							
Throwing things							
Spitting							
Tearing things/destroying property							
Grabbing onto people							
Other:							

Additional behaviors include making physical sexual advances, intentional falling, eating/drinking inappropriate substances, hiding things, hoarding things, making verbal sexual advances, and strange noises (weird laughter or crying).

otes:				
	1.0			
naviors nav	ve caused dis	ruption to:		





